

Contents

Editorial

- Epidural ketamine for postoperative analgesia/*Kétamine péridurale pour l'analgésie postopératoire*
A.N. Sandler, R. Schmid, J. Katz 99

Reports of Investigation

- Epidural ketamine reduces post-operative epidural PCA consumption of fentanyl/bupivacaine
M.E. Abdel-Ghaffar, M. Abdulatif, A. Al-Ghamdi, H. Mowafi, A. Anwar 103
- North American survey of the management of dural puncture occurring during labour epidural analgesia
C.W. Berger, E.T. Crosby, W. Grodecki 110
- 40 Hz Auditory steady-state response and EEG spectral edge frequency during sufentanil anaesthesia
I. Gilron, G. Plourde, W. Marcantoni, F. Varin 115
- Minimum effective anaesthetic concentration of hyperbaric lidocaine for spinal anaesthesia
P.W.H. Peng, V.W.S. Chan, A. Perlas 122
- Cardiac arrest in the OR: How are our ACLS skills?
M.M. Kurrek, J.H. Devitt, M. Cohen 130
- Cardiac output during liver transplantation: comparison between thermodilution and transoesophageal Doppler monitoring
S. Colbert, D.M. O'Hanlon, J. Duranteau, C. Ecoffey 133
- Haemodynamic effects of rocuronium bromide in adult cardiac surgical patients
M.E. Hudson, K.P. Rothfield, W.C. Tullock, L.L. Firestone 139

- Predictors of hospital mortality and mechanical ventilation in patients with cervical spinal cord injury
A.R. Claxton, D.T. Wong, F. Chung, M.G. Fehlings 144

- Awake removal of the laryngeal mask airway is safe in paediatric patients
A.H. Samarkandi 150

- Prevention of PONV with granisetron, droperidol or metoclopramide in patients with postoperative emesis
Y. Fujii, Y. Saitoh, H. Tanaka, H. Toyooka 153

Obstetric Forum

- Anaesthetic management of Caesarean section in a patient with Myelodysplastic Syndrome
K. Hara, Y. Saito, N. Morimoto, S. Sakura, Y. Kosaka 157

Clinical Reports

- Transcutaneous PCO₂ monitoring during laparoscopic cholecystectomy in pregnancy
K. Bhavani-Shankar, R.A. Steinbrook, P.S. Mushlin, D. Freiberger 164
- A surgically placed epidural catheter in a patient with spinal trauma
D.D. Seal, R.G. Loken, R.J. Hurlbert 170

Preliminary Report

- Long-term epidural ketamine, morphine and bupivacaine attenuate reflex sympathetic dystrophy neuralgia
T.-C. Lin, C.-S. Wong, F.-C. Chen, S.-Y. Lin, S.-T. Ho 175

