

Contents

Editorials

- Somatosensory evoked potentials during spinal surgery/Les potentiels évoqués somatosensitifs durant la chirurgie du rachis
R.A. McTaggart Cowan 387
- The best approach to ARDS?/Quelle est la meilleure approche au SDRA ?
G. Blaise, E. Troncy 393

Reports of Investigation

- A rapid precurarization technique using rocuronium
B.C.H. Tsui, S. Reid, S. Gupta, R. Kearney, T. Mayson, B. Finucane 397
- Combined nitric oxide inhalation, prone positioning and almitrine infusion improve oxygenation in severe ARDS
T. Gillart, J.E. Bazin, B. Cosserant, D. Guelon, L. Aigouy, O. Mansoor, P. Schoeffler 402
- Age related variability in the effects of mivacurium in paediatric surgical patients
B.W. Brandom, O.A. Meretoja, E. Simhi, T. Taivainen, S.R. Wolfe, S.K. Woelfel, B.J. Gronert, D.R. Cook 410
- Propofol reduces succinylcholine induced increase of masseter muscle tone
W.C. Ummenhofer, C. Kindler, G. Tschalèr, K.F. Hampl, J. Drewe, A. Urwyler 417
- Electro cardiographic monitoring in healthy young adult outpatients: mandatory or optional?
I. Zabani, H. Vaghadia, C.R. Chilvers, P.M. Merrick 424
- Comparative effects of desflurane and isoflurane on recovery after long lasting anaesthesia
M. Beaussier, H. Deriaz, Z. Abdelahim, F. Aissa, A. Lienhart 429

- Comparison of tramadol and morphine via subcutaneous PCA following major orthopaedic surgery
D. Hopkins, E.A. Shipton, D. Potgieter, C.A. Van Der Merwe, J. Boon, C. De Wet, J. Murphy 435

- Volume kinetics of Ringer's solution and dextran 3% during induction of spinal anaesthesia for Caesarean section
R.G. Hahn, M. Resby 443

Obstetric Forum

- Obstetrical anaesthesia for a parturient with preeclampsia, HELLP syndrome and acute cortical blindness
E.T. Crosby, R. Preston 452

Clinical Measurement

- Monitoring evoked potentials during spinal surgery in one institution
P.H. Manninen 460

Laboratory Investigations

- Nonlinear additivity of nitrous oxide and isoflurane potencies in rats
G.B. Russell, J.M. Graybeal 466
- Inhibition by diazepam of ketamine-induced hyperlocomotion and dopamine turnover in mice
M. Irifune, T. Sato, Y. Kamata, T. Nishikawa, M. Nomoto, T. Fukuda, M. Kawahara 471
- Lower oesophageal sphincter tone increases after induction of anaesthesia in pigs with full stomach
J.-P. Tournadre, M. Barclay, P. Boulétreau, D. Chassard 479

<i>D.J. Doyle</i>	498
Textbook of Neuroanesthesia with Neurosurgical and Neuroscience Perspectives MS. <i>Albin (Ed.)</i> REVIEWER: <i>D. Archer</i>	498
Drugs in Anaesthesia & Intensive Care, 2nd ed. <i>M. Sasada, S. Smith</i> REVIEWERS: <i>W.J. Perks, P. Houston</i>	499
The ICU Book, 2nd ed. <i>PL. Marino</i> REVIEWER: <i>/ . Hooper</i>	499
Non-Standard Medical Electives in the U.S. and Canada <i>K. V. Iseron</i> REVIEWER: <i>D. Biehl</i>	499